



**Phoenix, Arizona — 1998 (N = 240)**


**Figure A. Age of GISP participants, in years, 1998**




**Figure B. Race/ethnicity of GISP participants, 1998**



**Figure C. Percentage of GISP participants Identifying as men who have sex with men**




**Figure D. Reason for visit among GISP participants, 1998**



## Phoenix, Arizona — 1998 (N=240)

**Figure E. Previous episode of gonorrhea among GISP participants**




\*Data first collected in 1991.


\*\*Data first collected in 1992.

Note: Data points not shown when >30% data missing.

**Figure F. Treatment given to GISP participants, 1998**




**Figure G. Resistance to penicillin and tetracycline among GISP isolates, 1998**




Pen & Tet Suscep	PPNG	TRNG
PPNG/TRNG	PenR	TetR
CMRNG		

**Figure H. Ceftriaxone minimum inhibitory concentrations (MIC) among GISP isolates, 1998**




## Phoenix, Arizona — 1998 (N=240)

**Figure I. Cefixime minimum inhibitory concentrations (MIC) among GISP isolates, 1998**



**Figure J. Intermediate resistance and resistance to ciprofloxacin among GISP isolates**



Note: Susceptibility to ciprofloxacin first measured in 1990.

**Figure K. Ciprofloxacin minimum inhibitory concentrations (MIC) among GISP isolates, 1998**

